Notes on squeezed states in x-space representation

Alexander N. Korotkov

Department of Electrical and Computer Engineering, University of California, Riverside, California 92521 (Dated: September 25, 2018)

In these notes, we discuss squeezed states using the elementary quantum language based on one-dimensional Schrödinger equation. No operators are used. The language of quantum optics is mentioned only for a hint to solve a differential equation. Sections II and III (squeezed vacuum and pure squeezed states) can be used in a standard undergraduate course on quantum mechanics after discussion of a harmonic oscillator. Section IV discusses density matrix of mixed squeezed states in x-representation. These notes present explicit (therefore somewhat lengthy) step-by-step derivations. These notes are not intended for publication as a journal paper.

I. INTRODUCTION

In quantum optics [1-4], squeezed states are usually introduced via the squeezing operator and discussed in the phase space, with the formalism based on creation and annihilation operators. Here we discuss one-dimensional squeezed states in the *x*-representation, using the formalism based only on elementary quantum mechanics [5, 6].

II. SQUEEZED VACUUM

Let us consider a harmonic oscillator with mass m and frequency ω , so the potential energy is

$$V(x) = \frac{1}{2}m\omega^2 x^2.$$
 (1)

The ground state of this oscillator is [5, 6]

$$\psi_{\rm gr}(x,t) = \frac{e^{-\imath\omega t/2}}{\sqrt[4]{2\pi\sigma_{\rm gr}^2}} \exp\left(-\frac{x^2}{4\sigma_{\rm gr}^2}\right), \ \sigma_{\rm gr}^2 = \frac{\hbar}{2m\omega}, \quad (2)$$

where $\sigma_{\rm gr}^2$ is the *x*-space variance of the ground state probability distribution $|\psi_{\rm gr}(t)|^2$.

Suppose we have prepared an initial state $\psi(x,0) = \exp(-x^2/4D)/\text{Norm.}$ What is the further evolution $\psi(x,t)$ of this state? If $D = \sigma_{\text{gr}}^2$, then we have prepared the ground state and therefore it will not evolve (except for the trivial time-factor $e^{-i\omega t/2}$ due to the ground state energy $\hbar\omega/2$). If $D \neq \sigma_{\text{gr}}^2$, then this state is called a squeezed vacuum (especially if $D < \sigma_{\text{gr}}^2$). Our goal in this section is to find the explicit time-dependence $\psi(x,t)$ for the squeezed vacuum in x-space.

In quantum optics language, the time-dependence of the squeezed vacuum corresponds to the rotation of quadrature axes with frequency ω . From the picture of an ellipse representing the squeezed vacuum in optics, we know that $\psi(x,t)$ should be periodic with frequency 2ω (except for an accumulating phase). And we also know that a Gaussian state should remain Gaussian with rotation of axes. Also, since $\psi(x,0)$ is symmetric (even), this symmetry should remain in time. Therefore, the squeezed vacuum should be

$$\psi(x,t) = \frac{1}{\sqrt[4]{2\pi\sigma_{\rm gr}^2 A(t)}} \exp\left(-\frac{x^2[1+iB(t)]}{4\sigma_{\rm gr}^2 A(t)}\right) e^{-i\varphi(t)},$$
(3)

where A(t), B(t), and $\varphi(t)$ are real dimensionless functions of time, with A(t) and B(t) being 2ω -periodic (note that this state is already normalized). In particular, for the ground state A(t) = 1, B(t) = 0, and $\varphi(t) = \omega t/2$. Let us find A(t), B(t), and $\varphi(t)$ for the squeezed vacuum. To use the Schrödinger equation [5, 6],

$$i\hbar\frac{\partial\psi}{\partial t} = -\frac{\hbar^2}{2m}\frac{\partial^2\psi}{dx^2} + \frac{1}{2}m\omega^2 x^2\psi, \qquad (4)$$

we need to calculate $\partial \psi / \partial t$ and $\partial^2 \psi / dx^2$. This can be easily done for the wavefunction (3),

$$\frac{\partial\psi}{\partial t} = \left[-\frac{\dot{A}}{4A} + \frac{x^2(1+iB)\dot{A}}{4\sigma_{\rm gr}^2 A^2} - \frac{ix^2\dot{B}}{4\sigma_{\rm gr}^2 A} - i\dot{\varphi} \right]\psi, \quad (5)$$

$$\frac{\partial^2 \psi}{\partial x^2} = \left[\left(-\frac{x(1+iB)}{2\sigma_{\rm gr}^2 A} \right)^2 - \frac{1+iB}{2\sigma_{\rm gr}^2 A} \right] \psi. \tag{6}$$

Substituting these derivatives into the Schrödinger equation, we obtain

$$i\hbar \left[-\frac{\dot{A}}{4A} + \frac{x^2(1+iB)\dot{A}}{4\sigma_{\rm gr}^2 A^2} - \frac{ix^2\dot{B}}{4\sigma_{\rm gr}^2 A} - i\dot{\varphi} \right] = -\frac{\hbar^2}{2m} \left[\frac{x^2(1+iB)^2}{4\sigma_{\rm gr}^4 A^2} - \frac{1+iB}{2\sigma_{\rm gr}^2 A} \right] + \frac{1}{2}m\omega^2 x^2, \quad (7)$$

which is the same as

_

.

$$-\frac{A}{A} + \frac{x^{2}(1+iB)A}{\sigma_{\rm gr}^{2}A^{2}} - \frac{ix^{2}B}{\sigma_{\rm gr}^{2}A} - 4i\dot{\varphi} = = \frac{i\hbar}{2m} \left[\frac{x^{2}(1+iB)^{2}}{\sigma_{\rm gr}^{4}A^{2}} - 2\frac{1+iB}{\sigma_{\rm gr}^{2}A} \right] - \frac{2i}{\hbar}m\omega^{2}x^{2}.$$
 (8)

Since this relation is valid for any x, it gives us two equations: for x^2 -tems and for x^0 -terms. Let us start with the equation for x^2 -component (multiplied by $\sigma_{\rm gr}^2$),

$$\frac{(1+iB)\dot{A}}{A^2} - \frac{i\dot{B}}{A} = \frac{i\hbar(1+iB)^2}{2m\sigma_{\rm gr}^2 A^2} - \frac{2i}{\hbar}m\omega^2\sigma_{\rm gr}^2.$$
 (9)

Using $\sigma_{\rm gr}^2 = \hbar/(2m\omega)$ and multiplying all terms by A^2 , we can rewrite Eq. (9) as

$$(1+iB)\dot{A} - iA\dot{B} = i\omega(1+iB)^2 - i\omega A^2.$$
 (10)

Since A and B are real, this equation gives us two equations: for the real and imaginary components,

$$\dot{A} = -2\omega B,\tag{11}$$

$$B\dot{A} - A\dot{B} = \omega(1 - B^2 - A^2).$$
 (12)

These two equations are sufficient to find A(t) and B(t). We will discuss the solution a little later; before that, let us consider the x^0 -component of Eq. (8), which is

$$-\frac{\dot{A}}{A} - 4i\dot{\varphi} = -\frac{i\hbar}{m}\frac{1+iB}{\sigma_{\rm gr}^2 A} = \frac{2\omega}{A}\left(-i+B\right).$$
(13)

The real part of this equation gives $\dot{A} = -2\omega B$, which is the same as Eq. (11). The imaginary part gives

$$\dot{\varphi} = \frac{\omega}{2A},\tag{14}$$

from which we can find $\varphi(t)$ if we know A(t).

Thus, we only have to solve Eqs. (11) and (12). Since we know that both A(t) and B(t) should be periodic with frequency 2ω , it is quite simple to guess the solution:

$$A(t) = A_0 + \Delta A \cos(2\omega t + \phi_{sq}), \qquad (15)$$

$$B(t) = \Delta A \sin(2\omega t + \phi_{sq}), \qquad (16)$$

where ϕ_{sq} is the initial phase for oscillations of squeezing and $A_0 > \Delta A \ge 0$. It is easy to see that Eq. (11) is satisfied for any A_0 , ΔA , and ϕ_{sq} , while to satisfy Eq. (12) we need the condition

$$(A_0 + \Delta A)(A_0 - \Delta A) = 1, \qquad (17)$$

which is a familiar condition for the product of minimum and maximum quadrature variances of a squeezed state. Note that for the initial state discussed at the beginning, we need $\phi_{sq} = n\pi$ with an integer n. Also note that the squeezed vacuum (3) is a minimum-uncertainty state, $\sigma_x \sigma_p = \hbar/2$, only when $2\omega t + \phi_{sq} = n\pi$ with integer n, i.e., four times per oscillator period.

Thus, we have found the evolution of the squeezed vacuum (3) up to the overall phase $\varphi(t)$, which is not quite important and can be found via Eq. (14). It is interesting that there is an explicit solution,

$$\varphi(t) = \frac{\omega}{2} \int^{t} \frac{dt'}{A_0 + \Delta A \cos(2\omega t' + \phi_{\rm sq})} \tag{18}$$

$$= \frac{1}{2} \operatorname{atan} \left[(A_0 - \Delta A) \tan(\omega t + \frac{\phi_{\mathrm{sq}}}{2}) \right] + \operatorname{const}, (19)$$

where each discontinuity of the tangent should be compensated by an increase of the constant by $\pi/2$. Note that in deriving Eq. (19) we have used Eq. (17). It is easy to see that for each $\Delta t = \pi/\omega$ (half-period), the phase increases by $\Delta \varphi = \pi/2$, so that $\Delta \varphi/\Delta t = \omega/2$, which is the same value as for the ground state.

III. PURE SQUEEZED STATES

Now let us generalize the x-space evolution of the squeezed vacuum to the x-space evolution of an arbitrary (pure) squeezed state. Since we know that for a squeezed state the evolution of the center is decoupled from squeezing, it is natural to guess the following wavefunction for a squeezed state:

$$\psi(x,t) = \frac{1}{\sqrt[4]{2\pi\sigma_{\rm gr}^2 A}} \exp\left[-\frac{(x-x_{\rm c})^2(1+iB)}{4\sigma_{\rm gr}^2 A}\right] \\ \times \exp(ip_{\rm c}x/\hbar) e^{-i\varphi}, \qquad (20)$$

where A(t) and B(t) are the same functions of time as for the squeezed vacuum [given by Eqs. (15)–(17)], while the overall phase $\varphi(t)$ is now different, and the state center evolves classically,

$$x_{\rm c} = X_{\rm amp} \cos(\omega t + \phi_{\rm c}), \qquad (21)$$

$$p_{\rm c} = m\dot{x}_{\rm c} = -m\omega X_{\rm amp}\sin(\omega t + \phi_{\rm c}),$$
 (22)

where $X_{\rm amp}$ is the amplitude of classical oscillations of the center and the initial phase $\phi_{\rm c}$ is not related to the initial phase $\phi_{\rm sq}$ of squeezing oscillations. Let us show that the time-dependence of an arbitrary squeezed state given by Eqs. (20)–(22) and (15)–(17) satisfy the Schrödinger equation.

The derivatives of $\psi(x,t)$ given by Eq. (20) are

$$\frac{\partial \psi}{\partial t} = \left[-\frac{\dot{A}}{4A} + \frac{(x-x_{\rm c})^2(1+iB)\dot{A}}{4\sigma_{\rm gr}^2 A^2} - \frac{i(x-x_{\rm c})^2 \dot{B}}{4\sigma_{\rm gr}^2 A} + \frac{(x-x_{\rm c})\dot{x}_{\rm c}(1+iB)}{2\sigma_{\rm gr}^2 A} + \frac{i\dot{p}_{\rm c}x}{\hbar} - i\dot{\varphi} \right] \psi, \quad (23)$$

$$\frac{\partial^2 \psi}{\partial x^2} = \left[\left[-\frac{(x - x_{\rm c})(1 + iB)}{2\sigma_{\rm gr}^2 A} + \frac{ip_{\rm c}}{\hbar} \right]^2 - \frac{1 + iB}{2\sigma_{\rm gr}^2 A} \right] \psi, (24)$$

so the Schrödinger equation is [instead of Eq. (8)]

$$-\frac{\dot{A}}{A} + \frac{(x-x_{\rm c})^{2}(1+iB)\dot{A}}{\sigma_{\rm gr}^{2}A^{2}} - \frac{i(x-x_{\rm c})^{2}\dot{B}}{\sigma_{\rm gr}^{2}A} - 4i\dot{\varphi} + \frac{2(x-x_{\rm c})\dot{x}_{\rm c}(1+iB)}{\sigma_{\rm gr}^{2}A} + 4i\frac{\dot{p}_{\rm c}x}{\hbar} =$$
$$= \frac{i\hbar}{2m} \left[\frac{(x-x_{\rm c})^{2}(1+iB)^{2}}{\sigma_{\rm gr}^{4}A^{2}} - 2\frac{1+iB}{\sigma_{\rm gr}^{2}A} - 4\frac{p_{\rm c}^{2}}{\hbar^{2}} - 4i\frac{p_{\rm c}}{\hbar}\frac{(x-x_{\rm c})(1+iB)}{\sigma_{\rm gr}^{2}A} \right] - \frac{2i}{\hbar}m\omega^{2}x^{2}. \quad (25)$$

This equation now gives three equations: for x^2 , x^1 , and x^0 components. It is easy to check that the x^2 component still gives Eq. (9). Therefore, Eqs. (11)–(12) for A(t) and B(t) and correspondingly their solutions, Eqs. (15)–(17), are still valid for an arbitrary squeezed state. The x^1 -component of Eq. (25) multiplied by σ_{gr}^2 gives

$$\frac{-2x_{\rm c}(1+iB)\dot{A}}{A^2} + \frac{2ix_{\rm c}\dot{B}}{A} + \frac{2\dot{x}_{\rm c}(1+iB)}{A} + 4i\frac{\dot{p}_{\rm c}}{\hbar}\sigma_{\rm gr}^2 = = \frac{i\hbar}{2m}\frac{-2x_{\rm c}(1+iB)^2}{\sigma_{\rm gr}^2A^2} + \frac{2p_{\rm c}}{m}\frac{1+iB}{A}.$$
 (26)

Note that several terms here are the same as the terms in Eq. (9) multiplied by $-2x_c$. Subtracting Eq. (9) multiplied by $-2x_c$ from Eq. (26) and using $\sigma_{\rm gr}^2 = \hbar/2m\omega$, we obtain

$$\frac{2\dot{x}_{\rm c}(1+iB)}{A} + 2i\,\frac{\dot{p}_{\rm c}}{m\omega} = \frac{2p_{\rm c}}{m}\frac{1+iB}{A} - 2ix_{\rm c}\omega,\qquad(27)$$

which is obviously satisfied for x_c and p_c given by Eqs. (21) and (22). Thus, the x^1 -component of Eq. (25) is satisfied. Finally, the x^0 -component gives

$$-\frac{\dot{A}}{A} + \frac{x_{\rm c}^2(1+iB)\dot{A}}{\sigma_{\rm gr}^2 A^2} - \frac{ix_{\rm c}^2\dot{B}}{\sigma_{\rm gr}^2 A} - 4i\dot{\varphi} - \frac{2x_{\rm c}\dot{x}_{\rm c}(1+iB)}{\sigma_{\rm gr}^2 A} = \sup_{\substack{\rm ave}} \frac{i\hbar}{\sigma_{\rm gr}^2 A} - \frac{x_{\rm c}^2\dot{A}}{\sigma_{\rm gr}^2 A} - 4\frac{p_{\rm c}^2}{\hbar^2} + 4i\frac{p_{\rm c}}{\hbar}\frac{x_{\rm c}(1+iB)}{\sigma_{\rm gr}^2 A} - (28)^{\rm and}$$

The real part of this equation is

$$-\frac{\dot{A}}{A} + \frac{x_{\rm c}^2 \dot{A}}{\sigma_{\rm gr}^2 A^2} - \frac{2x_{\rm c} \dot{x}_{\rm c}}{\sigma_{\rm gr}^2 A} = \frac{\hbar B}{m\sigma_{\rm gr}^2 A} - \frac{\hbar x_{\rm c}^2 B}{m\sigma_{\rm gr}^4 A^2} - 2\frac{p_{\rm c} x_{\rm c}}{m\sigma_{\rm gr}^2 A}.$$
(29)

The first terms on both sides are equal to each other because $\dot{A} = -2\omega B$ [see Eq. (11)]. The second terms are equal because of the same reason. The third terms are equal because $\dot{x}_c = p_c/m$. So, Eq. (29) is satisfied. Finally, the imaginary part of Eq. (28) gives (after some algebra)

$$\dot{\varphi} = \frac{\omega}{2A} + \frac{x_{\rm c}^2 B \dot{A}}{4\sigma_{\rm gr}^2 A^2} - \frac{x_{\rm c}^2 \dot{B}}{4\sigma_{\rm gr}^2 A} - \frac{\omega x_{\rm c}^2 (1 - B^2)}{4\sigma_{\rm gr}^2 A^2} + \frac{p_{\rm c}^2}{2\hbar m} = \frac{\omega}{2A} + \frac{m\omega^2 (X_{\rm amp}^2 - x_{\rm c}^2)}{2\hbar}.$$
(30)

It is simple to solve this equation by adding the explicit solution (19) for $x_{\rm c} = X_{\rm amp} = 0$ and the integral of the last term in Eq. (30).

Thus, we have shown that the general (pure) squeezed state in x-space has the form given by Eqs. (20)-(22), (15)-(17), and (30).

IV. MIXED SQUEEZED (GAUSSIAN) STATES

Using the wavefunction (20), let us construct the density matrix

$$\begin{split}
\rho(x,x') &= \psi(x)\,\psi^*(x') = \frac{1}{\sqrt{2\pi\sigma_{\rm gr}^2 A}}\,\exp\left[-\frac{(\frac{x+x'}{2}-x_{\rm c})^2}{2\sigma_{\rm gr}^2 A}\right] \\
&\times \exp\left[-\frac{(\frac{x-x'}{2})^2}{2\sigma_{\rm gr}^2 A}\right]\,\exp\left[-iB\frac{(\frac{x+x'}{2}-x_{\rm c})(x-x')}{2\sigma_{\rm gr}^2 A}\right] \\
&\times \exp\left[i\frac{p_{\rm c}(x-x')}{\hbar}\right].
\end{split}$$
(31)

Now let us probabilistically mix the states with $x_{\rm c}$ and $p_{\rm c}$ having the Gaussian probability distribution

$$p(x_{\rm c}, p_{\rm c}) = \frac{1}{\sqrt{2\pi\sigma_{\rm a}^2}} \exp\left[-\frac{(x_{\rm c} - \bar{x}_{\rm c})^2}{2\sigma_{\rm a}^2}\right] \\ \times \frac{1}{\sqrt{2\pi(m\omega\sigma_{\rm a})^2}} \exp\left[-\frac{(p_{\rm c} - \bar{p}_{\rm c})^2}{2(m\omega\sigma_{\rm a})^2}\right], \quad (32)$$

so that the additional x_c -spread has variance σ_a^2 and the spread of $p_c/m\omega$ has the same variance σ_a^2 , while the averaged centers \bar{x}_c and \bar{p}_c evolve according to Eqs. (21) and (22), i.e.,

$$\dot{\bar{x}}_{\rm c} = \bar{p}_{\rm c}/m, \ \dot{\bar{p}}_{\rm c} = -m\omega^2 \bar{x}_{\rm c}.$$
(33)

It is easy to see that the evolution (33) of \bar{x}_c and \bar{p}_c does not change the probability distribution (32) for x_c and p_c , which evolve according to Eqs. (21)–(22). The easiest way to see this fact is to consider an evolution on the plane of x_c and $p_c/m\omega$, so that Eqs. (21)–(22) as well as (33) describe a rotation with frequency ω along a circle, while the Gaussian distribution (32) is isotropic. Therefore, if we average the density matrix (31) over x_c and p_c with the distribution (32), the result *automatically* satisfies the Schrödinger equation.

For averaging the density matrix (31) over x_c , the following formula is useful:

$$\int_{-\infty}^{\infty} \frac{\exp\left[-\frac{(x+y)^2 + a(x+y)}{2\sigma_1^2}\right]}{\sqrt{2\pi\sigma_1^2}} \frac{\exp\left(-\frac{y^2}{2\sigma_2^2}\right)}{\sqrt{2\pi\sigma_2^2}} \, dy = \\ = \frac{\exp\left[-\frac{x^2 + ax}{2(\sigma_1^2 + \sigma_2^2)}\right]}{\sqrt{2\pi(\sigma_1^2 + \sigma_2^2)}} \, \exp\left[\frac{a^2\sigma_2^2}{8\sigma_1^2(\sigma_1^2 + \sigma_2^2)}\right], \quad (34)$$

where a can be complex. Similarly, for averaging over $p_{\rm c}$ we can use

$$\int_{-\infty}^{\infty} e^{a(x+y)} \frac{\exp(-y^2/2\sigma^2)}{\sqrt{2\pi\sigma^2}} \, dy = e^{ax} \exp(a^2\sigma^2/2).$$
(35)

Therefore, averaging of the density matrix (31) over the

distribution (32) gives

$$\rho(x, x') = \frac{1}{\sqrt{2\pi\sigma_{\Sigma}^{2}}} \exp\left[-\frac{(\frac{x+x'}{2} - \bar{x}_{c})^{2}}{2\sigma_{\Sigma}^{2}}\right] \\ \times \exp\left[-iB\frac{(\frac{x+x'}{2} - \bar{x}_{c})(x-x')}{2\sigma_{\Sigma}^{2}}\right] \exp\left[i\frac{\bar{p}_{c}(x-x')}{\hbar}\right] \\ \times \exp\left[-\frac{(\frac{x-x'}{2})^{2}}{2\sigma_{gr}^{2}A} - \frac{B^{2}\sigma_{a}^{2}(x-x')^{2}}{8\sigma_{gr}^{2}A\sigma_{\Sigma}^{2}} - \frac{\sigma_{a}^{2}(x-x')^{2}}{8\sigma_{gr}^{4}}\right], (36)$$

where $\sigma_{\Sigma}^2 = \sigma_{\rm gr}^2 A + \sigma_{\rm a}^2$. Using Eqs. (15)–(17), the last exponential factor can be expressed as

$$\exp\left[-\frac{\left(\frac{x-x'}{2}\right)^2}{2\sigma_{\Sigma}^2} \frac{\sigma_{\rm gr}^4 + \sigma_{\rm a}^4 + 2A_0\sigma_{\rm gr}^2\sigma_{\rm a}^2}{\sigma_{\rm gr}^4}\right].$$
 (37)

We see that if we introduce $\tilde{A} \equiv A + \sigma_{\rm a}^2/\sigma_{\rm gr}^2$ and $\tilde{A}_0 \equiv A_0 + \sigma_{\rm a}^2/\sigma_{\rm gr}^2$, then $\sigma_{\Sigma}^2 = \sigma_{\rm gr}^2 \tilde{A}$ and the second fraction in Eq. (37) can be rewritten as $1 + (\sigma_{\rm a}/\sigma_{\rm gr})^4 + 2A_0(\sigma_{\rm a}/\sigma_{\rm gr})^2 = P = (\tilde{A}_0 + \Delta A)(\tilde{A}_0 - \Delta A)$.

Therefore, removing tilde signs and overbars $(\tilde{A} \to A, \tilde{A}_0 \to A_0, \bar{x}_c \to x_c, \bar{p}_c \to p_c)$, we can rewrite the averaged density matrix (36) as

$$\rho(x, x') = \frac{1}{\sqrt{2\pi\sigma_{\rm gr}^2 A}} \exp\left[-\frac{(\frac{x+x'}{2} - x_{\rm c})^2}{2\sigma_{\rm gr}^2 A}\right]$$
$$\times \exp\left[-P\frac{(\frac{x-x'}{2})^2}{2\sigma_{\rm gr}^2 A}\right] \exp\left[-iB\frac{(\frac{x+x'}{2} - x_{\rm c})(x-x')}{2\sigma_{\rm gr}^2 A}\right]$$
$$\times \exp\left[i\frac{p_{\rm c}(x-x')}{\hbar}\right], \qquad (38)$$

- [1] D. F. Walls and G. J. Milburn, *Quantum Optics* (Springer, 2008).
- [2] C. C. Gerry and P. Knight, *Introductory Quantum Optics* (Cambridge University Press, 2005).
- [3] M. O. Scully and M. S. Zubairy, *Quantum Optics* (Cambridge University Press, 1997).
- [4] C. Gardiner and P. Zoller, Quantum Noise (Springer,

which differs from Eq. (31) for the pure state by only the factor P in the second exponent. This is the general form of a Gaussian state of an oscillator. In Eq. (38), the time-dependence of the center (x_c, p_c) is classical and is given by Eqs. (21) and (22); the center oscillates with frequency ω . The parameters A(t) and B(t) are given by Eqs. (15) and (16), they oscillate with frequency 2ω . The parameter P is the product of maximum and minimum dimensionless variances,

$$P = A_{\max}A_{\min} = (A_0 + \Delta A)(A_0 - \Delta A) \ge 1,$$
 (39)

so that P = 1 corresponds to a pure state, while P > 1 corresponds to a mixed Gaussian state.

V. SUMMARY

In the x-space representation, the squeezed vacuum wavefunction is given by Eq. (3) with dimensionless variance A(t) and phase parameter B(t) given by Eqs. (15)–(17), and the overall phase $\varphi(t)$ given by Eq. (19). For an oscillator with frequency ω , the shape of the wavefunction oscillates with frequency 2ω .

The wavefunction of a pure squeezed state is given by Eq. (20) with the same parameters A(t) and B(t) as for the squeezed vacuum, while the center position $x_c(t)$ and momentum $p_c(t)$ are given by Eqs. (21) and (22). The center oscillates with frequency ω , while the variance oscillates with frequency 2ω .

The density matrix of a mixed (Gaussian) squeezed state is given by Eq. (38) with the same equations for A(t), B(t), $x_c(t)$, and $p_c(t)$, but with an additional parameter P given by Eq. (39), which replaces Eq. (17).

2004).

- [5] D. J. Griffiths, Introduction to quantum mechanics (Pearson Prentice Hall, 2005).
- [6] J. J. Sakurai, Modern quantum mechanics (Addison-Wesley, 1994).